

Compactness of the space of causal curves

Keye Martin

Department of Mathematics
Tulane University
New Orleans, LA 70118

`martin@math.tulane.edu` · <http://math.tulane.edu/~martin>

Abstract

We prove that the space of causal curves between compact subsets of a separable globally hyperbolic poset is itself compact in the Vietoris topology. Although this result implies the usual result in general relativity, its proof does not require the use of geometry or differentiable structure.

1 Introduction

An important result in general relativity is that the space of causal curves between two compact sets in a globally hyperbolic spacetime is itself compact in the Vietoris (and hence upper) topology. This result plays an important role in the proofs of certain singularity theorems [13], in establishing the existence of maximum length geodesics [3], and in the proof of certain positive mass theorems [10].

Recently [8] it was shown that a globally hyperbolic spacetime $\mathcal{M}$ with its causality relation forms a special type of bicontinuous poset (i.e., a *globally hyperbolic poset*) which has a canonical representation by a *domain* $\mathbf{IM}$. This implies for example that a globally hyperbolic spacetime can be order theoretically reconstructed from only a countable dense set of events and timelike causality. It also suggests that the natural way to topologize the space of causal curves in general relativity is with the Vietoris topology. The reason for this is as follows.

First, there is a homeomorphism $\mathcal{M} \simeq \max(\mathbf{IM})$ between the manifold topology and the relative *Scott* topology on $\max(\mathbf{IM})$. Thus, *events* are maximal elements in $\mathbf{IM}$. Next, given any ω -continuous domain D with

$\max(D)$ metrizable, there is a ‘higher dimensional’ domain $\mathbf{CD}$ called the *convex powerdomain* which admits an injection

$$\mathcal{P}_{com}(\max(D)) \rightarrow \max(\mathbf{CD})$$

of the nonempty compact subsets of $\max(D)$ (in its relative Scott topology) into the maximal elements of $\mathbf{C}(D)$. In particular, *causal curves* in a globally hyperbolic spacetime are maximal elements in $\mathbf{C}(\mathbf{IM})$. Just as the relative Scott topology on $\max(\mathbf{IM})$ gives the manifold topology, the relative Scott topology on $\max(\mathbf{C}(\mathbf{IM}))$, when restricted to the causal curves, is exactly the Vietoris topology! When coupled with the fact that compactness in the Vietoris topology implies compactness in the upper topology, the use of the Vietoris topology in [12] is definitely natural, and probably aesthetically necessary.

So as stated above we prove that the space of causal curves between compact sets in a globally hyperbolic poset is compact in the Vietoris topology. The fact that the proof is entirely order theoretic seems to provide evidence that globally hyperbolic posets provide an abstract formulation of the physical notion causality which is independent of geometry and differentiable structure.

2 Domains

A *poset* is a partially ordered set, i.e., a set together with a reflexive, anti-symmetric and transitive relation.

Definition 2.1 Let $(P, \sqsubseteq)$ be a partially ordered set. A nonempty subset $S \subseteq P$ is *directed* if $(\forall x, y \in S)(\exists z \in S) x, y \sqsubseteq z$. The *supremum* of $S \subseteq P$ is the least of all its upper bounds provided it exists. This is written $\bigsqcup S$.

These ideas have duals that will be important to us: A nonempty $S \subseteq P$ is *filtered* if $(\forall x, y \in S)(\exists z \in S) z \sqsubseteq x, y$. The *infimum* $\bigwedge S$ of $S \subseteq P$ is the greatest of all its lower bounds provided it exists.

Definition 2.2 For a subset X of a poset P , set

$$\uparrow X := \{y \in P : (\exists x \in X) x \sqsubseteq y\} \quad \& \quad \downarrow X := \{y \in P : (\exists x \in X) y \sqsubseteq x\}.$$

We write $\uparrow x = \uparrow \{x\}$ and $\downarrow x = \downarrow \{x\}$ for elements $x \in X$.

A partial order allows for the derivation of several intrinsically defined topologies. Here is our first example.

Definition 2.3 A subset U of a poset P is *Scott open* if

- (i) U is an upper set: $x \in U \ \& \ x \sqsubseteq y \Rightarrow y \in U$, and
- (ii) U is inaccessible by directed suprema: For every directed $S \subseteq P$ with a supremum,

$$\bigsqcup S \in U \Rightarrow S \cap U \neq \emptyset.$$

The collection of all Scott open sets on P is called the *Scott topology*.

Definition 2.4 A *dcpo* is a poset in which every directed subset has a supremum. The *least element* in a poset, when it exists, is the unique element $\perp$ with $\perp \sqsubseteq x$ for all x .

The set of *maximal elements* in a dcpo D is

$$\max(D) := \{x \in D : \uparrow x = \{x\}\}.$$

Each element in a dcpo has a maximal element above it.

Definition 2.5 For elements x, y of a poset, write $x \ll y$ iff for all directed sets S with a supremum,

$$y \sqsubseteq \bigsqcup S \Rightarrow (\exists s \in S) \ x \sqsubseteq s.$$

We set $\downarrow x = \{a \in D : a \ll x\}$ and $\uparrow x = \{a \in D : x \ll a\}$.

For the symbol “ $\ll$,” read “approximates.”

Definition 2.6 A *basis* for a poset D is a subset B such that $B \cap \downarrow x$ contains a directed set with supremum x for all $x \in D$. A poset is *continuous* if it has a basis. A poset is ω -*continuous* if it has a countable basis.

Continuous posets have an important property, they are *interpolative*.

Proposition 2.7 If $x \ll y$ in a continuous poset P , then there is $z \in P$ with $x \ll z \ll y$.

This enables a clear description of the Scott topology,

Theorem 2.8 The collection $\{\uparrow x : x \in D\}$ is a basis for the Scott topology on a continuous poset.

And also helps us give a clear definition of the *Lawson topology*.

Definition 2.9 The *Lawson topology* on a continuous poset P has as a basis all sets of the form $\uparrow x \setminus \uparrow F$, for $F \subseteq P$ finite.

Definition 2.10 A *continuous dcpo* is a continuous poset which is also a dcpo. A *domain* is a continuous dcpo.

We now consider some examples that illustrate the basic ideas.

Example 2.11 Let X be a locally compact Hausdorff space. Its *upper space*

$$\mathbf{U}X = \{\emptyset \neq K \subseteq X : K \text{ is compact}\}$$

ordered under reverse inclusion

$$A \sqsubseteq B \Leftrightarrow B \subseteq A$$

is a continuous dcpo:

- For directed $S \subseteq \mathbf{U}X$, $\bigsqcup S = \bigcap S$.
- For all $K, L \in \mathbf{U}X$, $K \ll L \Leftrightarrow L \subseteq \text{int}(K)$.
- $\mathbf{U}X$ is ω -continuous iff X has a countable basis.

It is interesting here that the space X can be recovered from $\mathbf{U}X$ in a purely order theoretic manner:

$$X \simeq \max(\mathbf{U}X) = \{\{x\} : x \in X\}$$

where $\max(\mathbf{U}X)$ carries the relative Scott topology it inherits as a subset of $\mathbf{U}X$. Several constructions of this type are known.

The next example is due to Scott[11]; it is good to understand it in detail, especially since globally hyperbolic spacetimes admit a completely analogous construction.

Example 2.12 The collection of compact intervals of the real line

$$\mathbf{I}\mathbb{R} = \{[a, b] : a, b \in \mathbb{R} \text{ \& } a \leq b\}$$

ordered under reverse inclusion

$$[a, b] \sqsubseteq [c, d] \Leftrightarrow [c, d] \subseteq [a, b]$$

is an ω -continuous dcpo:

- For directed $S \subseteq \mathbf{IR}$, $\bigsqcup S = \bigcap S$,
- $I \ll J \Leftrightarrow J \subseteq \text{int}(I)$, and
- $\{[p, q] : p, q \in \mathbb{Q} \text{ \& } p \leq q\}$ is a countable basis for $\mathbf{IR}$.

The domain $\mathbf{IR}$ is called the *interval domain*.

We also have $\max(\mathbf{IR}) \simeq \mathbb{R}$ in the Scott topology. The reason for this is that a basic Scott open set in $\mathbf{IR}$ has the form

$$\uparrow[a, b] = \{x \in \mathbf{IR} : x \subseteq (a, b)\}$$

so when the Scott topology is restricted to $\max(\mathbf{IR})$, we get open sets of the form

$$\uparrow[a, b] \cap \max(\mathbf{IR}) = \{[x] : x \in (a, b)\} \simeq (a, b)$$

which is just the Euclidean topology on $\mathbb{R}$.

3 Globally hyperbolic posets

In this section we consider a very special example of a domain. But first:

Definition 3.1 A continuous poset P is *bicontinuous* if

- For all $x, y \in P$, $x \ll y$ iff for all filtered $S \subseteq P$ with an infimum,

$$\bigwedge S \subseteq x \Rightarrow (\exists s \in S) s \subseteq y,$$

and

- For each $x \in P$, the set $\uparrow x$ is filtered with infimum x .

Example 3.2 $\mathbb{R}, \mathbb{Q}$ are bicontinuous.

Definition 3.3 On a bicontinuous poset P , sets of the form

$$(a, b) := \{x \in P : a \ll x \ll b\}$$

form a basis for a topology called *the interval topology*.

That the open intervals form a basis for a topology uses interpolation and bicontinuity. On a bicontinuous poset, the Lawson topology is contained in the interval topology (causal simplicity), the interval topology is Hausdorff (strong causality), and $\leq$ is a closed subset of P^2 . Globally hyperbolic posets provide a special example of a domain.

Definition 3.4 A poset $(X, \leq)$ is *globally hyperbolic* if it is bicontinuous and if the sets

$$[a, b] := \{x \in X : a \leq x \leq b\}$$

are compact in the interval topology.

Theorem 3.5 *The closed intervals of a globally hyperbolic poset X*

$$\mathbf{I}X := \{[a, b] : a \leq b \text{ \& } a, b \in X\}$$

ordered by reverse inclusion

$$[a, b] \sqsupseteq [c, d] \equiv [c, d] \subseteq [a, b]$$

form a continuous domain with

$$[a, b] \ll [c, d] \equiv a \ll c \text{ \& } d \ll b.$$

X has a countable basis iff $\mathbf{I}X$ is ω -continuous. Finally,

$$\max(\mathbf{I}X) \simeq X$$

where the set of maximal elements has the relative Scott topology from $\mathbf{I}X$, and X has the interval topology.

In fact there is even an equivalence between globally hyperbolic posets and certain types of domains [8]. We briefly now recall the relevance of these ideas to general relativity; more detailed definitions are in [8].

Definition 3.6 A *spacetime* is a real four-dimensional smooth manifold $\mathcal{M}$ with a Lorentz metric g_{ab} .

Let $(\mathcal{M}, g_{ab})$ be a time orientable spacetime. Let $\Pi_{\leq}^+$ denote the future directed causal curves, and $\Pi_{<}^+$ denote the future directed time-like curves.

Definition 3.7 For $p \in \mathcal{M}$,

$$I^+(p) := \{q \in \mathcal{M} : (\exists \pi \in \Pi_{<}^+) \pi(0) = p, \pi(1) = q\}$$

and

$$J^+(p) := \{q \in \mathcal{M} : (\exists \pi \in \Pi_{\leq}^+) \pi(0) = p, \pi(1) = q\}$$

Similarly, we define $I^-(p)$ and $J^-(p)$.

We write the relation J^+ as

$$p \leq q \equiv q \in J^+(p).$$

Definition 3.8 A spacetime $\mathcal{M}$ is *globally hyperbolic* if it is strongly causal and if $\uparrow a \cap \downarrow b$ is compact in the manifold topology, for all $a, b \in \mathcal{M}$.

Theorem 3.9 *If $(\mathcal{M}, \leq)$ is globally hyperbolic, then $(\mathcal{M}, \sqsubseteq)$ is a bicontinuous poset with $\ll = I^+$ whose interval topology is the manifold topology.*

Thus, a globally hyperbolic spacetime is a globally hyperbolic poset when equipped with its causality relation $\leq$.

4 The space of causal curves

A fundamental result in relativity is that the space of causal curves between points is compact on a globally hyperbolic spacetime. We use domains as an aid in proving this fact for any globally hyperbolic poset (they are not necessary though). One advantage to involving domains in the picture is that the Vietoris topology on causal curves arises as the natural counterpart to the manifold topology on events, so we can understand that its use in [12] is very natural.

In addition, other results on the compactness of the space of causal curves, such as those used to establish the existence of maximum length geodesics, are easily derivable from this one result. In fact, the length function is continuous with respect to the Vietoris topology as well, so it is possible to make a very strong case that the Vietoris topology is more natural than the one normally used (the “upper” topology).

Definition 4.1 Let D be a continuous dcpo. For subsets $A, B \subseteq D$, we define relations

- $A \ll_L B \Leftrightarrow (\forall a \in A)(\exists b \in B) a \ll b$
- $A \ll_U B \Leftrightarrow (\forall b \in B)(\exists a \in A) a \ll b$
- $A \ll_{EM} B \Leftrightarrow A \ll_L B \ \& \ A \ll_U B$

In the same way, we derive $\sqsubseteq_L, \sqsubseteq_U$ and $\sqsubseteq_{EM}$ from the order $\sqsubseteq$ on D .

Definition 4.2 The nonempty finite subsets of a space X are denoted $\mathcal{P}_{fin}(X)$, while its nonempty compact subsets are written as $\mathcal{P}_{com}(X)$.

The set $\mathcal{P}_{fin}(D)$ together with $\ll_{EM}$ is an abstract basis. (See the appendix for more on abstract bases).

Definition 4.3 The *convex powerdomain* $\mathbf{CD}$ of a continuous dcpo D is the ideal completion of the abstract basis $(\mathcal{P}_{fin}(D), \ll_{EM})$.

Definition 4.4 For a Scott compact $K \in \mathcal{P}_{com}(D)$, we set

$$K^* = \{F \in \mathcal{P}_{fin}(D) : F \ll_{EM} K\}.$$

Notice that this operation is also defined for elements of $\mathcal{P}_{fin}(D)$.

Proposition 4.5 For a continuous dcpo D , we have

- (i) If $K \in \mathcal{P}_{com}(D)$, then $K^* = \{F \in \mathcal{P}_{fin}(D) : F \ll_{EM} K\} \in \mathbf{CD}$.
- (ii) For ideals $I, J \in \mathbf{CD}$,

$$I \ll J \Leftrightarrow (\exists F, G \in \mathcal{P}_{fin}(D)) F \ll_{EM} G \text{ \& } I \subseteq F^* \subseteq G^* \subseteq J.$$

- (iii) For $F \in \mathcal{P}_{fin}(D)$ and $I \in \mathbf{CD}$, $F \in I \Leftrightarrow F^* \ll I$.
- (iv) For $F, G \in \mathcal{P}_{fin}(D)$, $F^* \subseteq G^*$ in $\mathbf{CD} \Leftrightarrow F \sqsubseteq_{EM} G$.

Definition 4.6 The *Vietoris hyperspace* of a Hausdorff space X is the set of all nonempty compact subsets $\mathcal{P}_{com}(X)$ with the *Vietoris topology*: It has a basis given by all sets of the form

$$\sigma(U_1, \dots, U_n) := \{K \in \mathcal{P}_{com}(X) : K \subseteq \bigcup_{i=1}^n U_i \text{ and } K \cap U_i \neq \emptyset, 1 \leq i \leq n\},$$

where U_i is a nonempty open subset of X , for each $1 \leq i \leq n$.

The next result is from [5]:

Theorem 4.7 Let D be an ω -continuous dcpo with $X = \max(D)$ regular in its relative Scott topology. Then the correspondence

$$\mathcal{P}_{com}(X) \xrightarrow{e} \max(\mathbf{CD}) :: K \mapsto K^*$$

is a homeomorphism between the Vietoris hyperspace of X and $\text{Im}(e)$ in its relative Scott topology.

Thus, events in spacetime are maximal elements in $\mathbf{IX}$, causal curves are maximal elements in $\mathbf{C}(\mathbf{IX})$. The reason we know this is:

Proposition 4.8 *A subset of a globally hyperbolic spacetime $\mathcal{M}$ is the image of a causal curve iff it is the image of a continuous monotone increasing $\pi : [0, 1] \rightarrow \mathcal{M}$ iff it is a compact connected linearly ordered subset of $(\mathcal{M}, \sqsubseteq)$.*

This suggests the following:

Definition 4.9 Let $(X, \leq)$ be a globally hyperbolic poset. A subset $\pi \subseteq X$ is a *causal curve* if it is compact, connected and linearly ordered. We define

$$\pi(0) := \perp \quad \text{and} \quad \pi(1) := \top$$

where $\perp$ and $\top$ are the least and greatest elements of π . For $P, Q \subseteq X$,

$$C(P, Q) := \{\pi : \pi \text{ causal curve}, \pi(0) \in P, \pi(1) \in Q\}$$

and call this *the space of causal curves* between P and Q .

Let D be an ω -continuous dcpo with $\max(D)$ regular. For example, it could be $D = \mathbf{IX}$ for X globally hyperbolic and separable.

Lemma 4.10 *If $\uparrow x \cap \max(D)$ is Scott compact in D , then $\uparrow\{x\}^* \cap \text{Im}(e)$ is Scott compact in $\mathbf{CD}$.*

Proof. Let $L := (\uparrow x \cap \max(D))$. It is a compact metric space. If $K^* \in \uparrow\{x\}^* \cap \text{Im}(e)$, then $\{x\}^* \subseteq K^*$, which implies $K \subseteq L$. Thus, $\uparrow\{x\}^* \cap \text{Im}(e) \simeq \mathcal{P}_{com}(L)$ in the Vietoris topology, which is also a compact metric space [2]. $\square$

For the next lemma, it is important to point out that $X = \max(D)$ is metrizable and thus *normal*.

Lemma 4.11 *If (k_i) is a convergent sequence in $\mathcal{P}_{com}(X)$ with each k_i connected, then $\lim k_i$ is connected.*

Proof. Assume $k := \lim k_i$ is disconnected. Then $k = a \cup b$ is a disjoint union of closed sets. Since k is compact, a and b are compact, so $a^*, b^* \in \text{Im}(e)$. By normality of X , there are disjoint open sets $U, V \subseteq X$ with $a \subseteq U$, $b \subseteq V$. Thus, there are finite sets $F, G \in \mathcal{P}_{fin}(D)$ with $a \subseteq \uparrow F \cap X \subseteq U$ and

$b \subseteq \uparrow G \cap X \subseteq V$. If any element of F or G is not way below some element of a or b , we can simply throw it out. Thus,

$$F^* \ll a^* \text{ and } G^* \ll b^* \Rightarrow (F \cup G)^* \ll (a \cup b)^* = k^*$$

so for large enough i , we have $(F \cup G)^* \ll (k_i)^*$, since $(k_i)^* \rightarrow k^*$ in the relative Scott topology. The sets $\uparrow F \cap k_i \cap X$ and $\uparrow G \cap k_i \cap X$ are open, disjoint and their union is k_i . They are nonempty because $F \cup G \ll_{EM} k_i$. Then k_i is disconnected. $\square$

In the next lemma, $D = \mathbf{IX}$ the domain of spacetime intervals for a separable globally hyperbolic poset.

Lemma 4.12 *If (k_i) is a convergent sequence in $\mathcal{P}_{com}(X)$ with each k_i linearly ordered, then $\lim k_i$ is linearly ordered.*

Proof. Let $k := \lim k_i$ and $p, q \in k$. There are increasing sequences (x_n) and (y_n) in $\mathbf{IX}$ with $x_n \ll [p]$, $y_n \ll [q]$, $\sqcup x_n = [p]$ and $\sqcup y_n = [q]$. By compactness of k , extend $\{x_n, y_n\}$ to a finite $F_n \supseteq \{x_n, y_n\}$ with $F_n^* \ll k^*$. Then

$$(\forall n)(\exists i_n)(k_{i_n})^* \in \uparrow F_n^*$$

Since $x_n, y_n \in F_n \ll_{EM} k_{i_n}$, there are $p_n, q_n \in k_{i_n}$ with $x_n \ll [p_n]$, $y_n \ll [q_n]$. Then $p_n \rightarrow p$ and $q_n \rightarrow q$. Because k_{i_n} is linearly ordered, we either have $p_n \leq q_n$ for an infinite number of n or $q_n \leq p_n$ for an infinite number of n . Because $\leq$ is closed, either $p \leq q$ or $q \leq p$, so k is linearly ordered. $\square$

In our final lemma, $(X, \leq)$ is a separable globally hyperbolic poset.

Lemma 4.13 *Let (π_n) be a sequence of causal curves with $\pi_n \rightarrow \pi$ in the Vietoris topology on $\mathcal{P}_{com}(X)$. If $(\pi_n(0))$ and $(\pi_n(1))$ are both convergent, then $\pi_n(0) \rightarrow \pi(0)$ and $\pi_n(1) \rightarrow \pi(1)$.*

Proof. First, we prove that for every $r \in \pi$, there is a subsequence $\alpha_n = \pi_{k_n}$ such that $r_n \rightarrow r$ where $r_n \in \alpha_n$. Take an increasing sequence (x_n) in $\mathbf{IX}$ with $x_n \ll [r]$ and $\sqcup x_n = [r]$. Extend $\{x_n\}$ to a finite set $F_n \supseteq \{x_n\}$ with $F_n \ll_{EM} \pi$ and thus $F_n^* \ll \pi^*$. For each n , let k_n be the least integer such that $F_n^* \ll \pi_{k_n}^*$ and hence $F_n \ll_{EM} \pi_{k_n}$. Since $x_n \in F_n$, there is $r_n \in \pi_{k_n}$ with $x_n \ll r_n$. But $\sqcup x_n = [r]$, $r_n \rightarrow r$, so $\alpha_n := \pi_{k_n}$ is the subsequence.

Next, we claim $a := \lim \pi_n(0)$ and $b := \lim \pi_n(1)$ belong to π . If $a \notin \pi$, then π and $\{a\}$ are disjoint compact sets, so by normality there are disjoint

open sets $U, V \subseteq X$ with $\pi \subseteq U$ and $a \in V$. But, since $\pi_n(0) \rightarrow a$, $\pi_n(0) \in V$ for all $n \geq L$, for some L . And, since $\pi_n \rightarrow \pi$ in the Vietoris topology, $\pi_n \subseteq U$ for all $n \geq L$, for some L . Thus, for n sufficiently large, $\pi_n(0) \in U \cap V$, contradicting $U \cap V = \emptyset$. A similar argument shows $b \in \pi$.

Finally, we can prove the claim. Given $r \in \pi$, take a subsequence (α_n) of (π_n) with $r_n \rightarrow r$ and $r_n \in \alpha_n$. Since (α_n) is a subsequence of (π_n) , each α_n is linearly ordered, so $\alpha_n(0) \leq r_n \leq \alpha_n(1)$. Then since $\leq$ is closed,

$$\lim_{n \rightarrow \infty} \alpha_n(0) = \lim_{n \rightarrow \infty} \pi_n(0) \leq r \leq \lim_{n \rightarrow \infty} \alpha_n(1) = \lim_{n \rightarrow \infty} \pi_n(1).$$

But $\lim \pi_n(0)$ and $\lim \pi_n(1)$ both belong to π and π is *linearly ordered*, so $\lim \pi_n(0) = \pi(0)$ and $\lim \pi_n(1) = \pi(1)$ since r was *arbitrary*. $\square$

Theorem 4.14 *If $(X, \leq)$ is a separable globally hyperbolic poset, then the space of causal curves $C(P, Q)$ is compact in the Vietoris topology when $P, Q \subseteq X$ are compact.*

Proof. Let (x_n) be a sequence in $C(P, Q)$. The endpoints $(x_n(0))$ have a convergent subsequence in P , so we pass to a subsequence of (x_n) called (y_n) with $(y_n(0))$ convergent. Then $(y_n(1))$ has a convergent subsequence in Q , so we pass to a subsequence of (y_n) called (π_n) which has the property that both $(\pi_n(0))$ and $(\pi_n(1))$ are convergent.

Define $a = \lim \pi_n(0) \in P$ and $b = \lim \pi_n(1) \in Q$. By definition, we have $\pi_n(0) \leq \pi_n(1)$ and thus $a \leq b$ since $\leq$ is closed. Then $[a, b] \in \mathbf{IX}$ so take some $u \ll [a, b]$ and notice that $\uparrow u \cap \max(\mathbf{IX})$ is an open set around $[a], [b]$. Since $\pi_n(0) \rightarrow a$ and $\pi_n(1) \rightarrow b$, we can assume $\pi_n(0), \pi_n(1) \in u$ for all n . But this implies $\pi_n \subseteq u$ because u is an interval! By global hyperbolicity, $\uparrow u \cap \max(\mathbf{IX})$ is compact, so $\uparrow \{u\}^* \cap \text{Im}(e)$ is compact in $\mathbf{C}(\mathbf{IX})$. Then (π_n) has a convergent subsequence in $\mathcal{P}_{com}(u)$ which we simply call (π_n) . Thus, $\pi_n \rightarrow \pi \in \mathcal{P}_{com}(u)$.

By our previous work, π is compact, connected and linearly ordered as a subset of $(X, \leq)$, so π is a causal curve. We still need to prove $\pi \in C(P, Q)$. By Lemma 4.13,

$$\lim \pi_n(0) = \pi(0) \text{ \& \; } \lim \pi_n(1) = \pi(1)$$

but we know these limits are exactly $a \in P$ and $b \in Q$. This proves $\pi \in C(P, Q)$. Since every sequence has a convergent subsequence and since $C(P, Q) \subseteq \mathcal{P}_{com}(X)$ has a countable basis (inherited from X), this proves $C(P, Q)$ is compact. $\square$

The separability is probably not required to prove this result, but we don't feel like working with nets anymore. The result above has been applied in the proof of a positive mass theorem [10]. We can also use it to deduce an important corollary, which is often used to establish the existence of maximum length geodesics on globally hyperbolic spacetimes ([3][13]).

Definition 4.15 The *upper topology* on $\mathcal{P}_{com}(X)$ has as a basis sets of the form

$$O(U) := \{K \in \mathcal{P}_{com}(X) : K \subseteq U\}$$

where $U \subseteq X$ is open.

Corollary 4.16 *If $(X, \leq)$ is a globally hyperbolic poset, the space of causal curves $C(P, Q)$ is compact in the upper topology whenever $P, Q \subseteq X$ are compact.*

Proof. The upper topology is contained in the Vietoris topology. $\square$

A spacetime is globally hyperbolic iff either of the space of causal curves is compact; the nontrivial direction is the one we have abstracted here to the level of posets. For separable globally hyperbolic posets which ‘mimic’ spacetime we can also prove the following equivalence:

Proposition 4.17 *Let $(X, \leq)$ be a separable locally compact bicontinuous poset such that $x \leq y$ iff there is a continuous monotone increasing $\pi : [0, 1] \rightarrow X$ with $\pi(0) = x$ and $\pi(1) = y$. The following are equivalent:*

- (i) *The poset $(X, \leq)$ is globally hyperbolic.*
- (ii) *The space $C(P, Q)$ is compact in the Vietoris topology when $P, Q \subseteq X$ are compact.*
- (iii) *The space $C(P, Q)$ is compact in the upper topology when $P, Q \subseteq X$ are compact.*

Proof. To prove (iii) $\Rightarrow$ (i), let $r_n \in [p, q]$ be a sequence. Then there is $\alpha_n : [0, 1/2] \rightarrow X$ from p to r_n and $\beta_n : [1/2, 1] \rightarrow X$ from r_n to q . We can paste them together to get $\pi_n : [0, 1] \rightarrow X$ from p to q . Because π_n is continuous and monotone, its image $\text{Im}(\pi_n)$ is a causal curve. Since $C(\{p\}, \{q\})$ is compact in the upper topology, $(\text{Im}(\pi_n))$ has a subsequence named (α_n) with $\alpha_n \rightarrow \alpha$. By local compactness, there is an open set $U \subseteq X$ with $\text{Cl}(U)$ compact and $\alpha \subseteq U$. Then there is an integer N such

that $r_{k_n} \in \alpha_n \subseteq \text{Cl}(U)$ for all $n \geq N$. By compactness, the sequence (r_{k_n}) has a convergent subsequence with limit r . But $r_{k_n} \in \alpha_n \subseteq [p, q]$ and $[p, q]$ is closed as the intersection of Lawson closed sets, so $r \in [p, q]$.

This proves (r_n) has a convergent subsequence in $[p, q]$ which by second countability proves compactness. $\square$

The property required of $(X, \leq)$ is interesting because it is one that domains in quantum mechanics also satisfy [7]. In closing this section, the fact that the compactness result can be proven for globally hyperbolic posets provides evidence that we have identified a useful order theoretic formulation of causality that does not use geometry or differentiable structure.

5 Conclusion

Sorkin and Woolgars in [12] recast the tools of “global causal analysis” using only topology and order. Here we have gone a step further, showing that only order is necessary, the topology is implicitly described by the order at an abstract level. Our results apply even to posets that are not spacetimes, demonstrating that in no way can our proof depend on geometry. However, we owe a great debt to the paper [12], which in our mind very clearly pointed the way to the proof of our main result.

References

- [1] S. Abramsky and A. Jung. *Domain theory*. In S. Abramsky, D. M. Gabbay, T. S. E. Maibaum, editors, *Handbook of Logic in Computer Science*, vol. III. Oxford University Press, 1994.
- [2] R. Engelking. *General topology*. Polish Scientific Publishers, Warszawa, 1977.
- [3] S.W. Hawking and G.F.R. Ellis. *The large scale structure of space-time*. Cambridge Monographs on Mathematical Physics, Cambridge University Press, 1973.
- [4] K. Martin. *A foundation for computation*. Ph.D. Thesis, Department of Mathematics, Tulane University, 2000.
- [5] K. Martin. *Fractals and domain theory*. Mathematical Structures in Computer Science, Cambridge University Press, to appear.

- [6] K. Martin. *The measurement process in domain theory*. Proc. 27th International Colloquium on Automata, Languages and Programming (ICALP), Lecture Notes in Computer Science, Vol. 1853, Springer-Verlag, 2000.
- [7] K. Martin. *Wronski brackets and the ferris wheel*. Research Report, Oxford University Computing Laboratory, April 2004.
- [8] K. Martin and P. Panangaden. *A domain of spacetime intervals in general relativity*. 2004.
- [9] R. Penrose. *Techniques of differential topology in relativity*. Society for Industrial and Applied Mathematics, 1972.
- [10] R. Penrose, R.D. Sorkin and E. Woolgar. *A positive mass theorem based on the focusing and retardation of null geodesics*. Submitted to Commun. Math. Phys., arXiv:gr-qc/9301015, 1993.
- [11] D. Scott. *Outline of a mathematical theory of computation*. Technical Monograph PRG-2, Oxford University Computing Laboratory, November 1970.
- [12] R.D. Sorkin and E. Woolgar. *A causal order for spacetimes with C^0 Lorentzian metrics: Proof of compactness of the space of causal curves*. Classical and Quantum Gravity **13**, p. 1971–1994, 1996.
- [13] R.M. Wald. *General relativity*. The University of Chicago Press, 1984.

Appendix: Abstract bases

A useful technique for constructing domains is to take the *ideal completion* of an *abstract basis*.

Definition 5.1 An *abstract basis* is given by a set B together with a transitive relation $<$ on B which is *interpolative*, that is,

$$M < x \Rightarrow (\exists y \in B) M < y < x$$

for all $x \in B$ and all finite subsets M of B .

Notice the meaning of $M < x$: It means $y < x$ for all $y \in M$. Abstract bases are covered in [1], which is where one finds the following.

Definition 5.2 An *ideal* in $(B, <)$ is a nonempty subset I of B such that

- (i) I is a lower set: $(\forall x \in B)(\forall y \in I) x < y \Rightarrow x \in I$.
- (ii) I is directed: $(\forall x, y \in I)(\exists z \in I) x, y < z$.

The collection of ideals of an abstract basis $(B, <)$ ordered under inclusion is a partially ordered set called the *ideal completion* of B . We denote this poset by $\bar{B}$.

The set $\{y \in B : y < x\}$ for $x \in B$ is an ideal which leads to a natural mapping from B into $\bar{B}$, given by $i(x) = \{y \in B : y < x\}$.

Proposition 5.3 If $(B, <)$ is an abstract basis, then

- (i) Its ideal completion $\bar{B}$ is a dcpo.
- (ii) For $I, J \in \bar{B}$,

$$I \ll J \Leftrightarrow (\exists x, y \in B) x < y \ \& \ I \subseteq i(x) \subseteq i(y) \subseteq J.$$

- (iii) $\bar{B}$ is a continuous dcpo with basis $i(B)$.

If one takes any basis B of a domain D and restricts the approximation relation $\ll$ on D to B , they are left with an abstract basis $(B, \ll)$ whose ideal completion is D . Thus, all domains arise as the ideal completion of an abstract basis.